

Table 3C.2 Reliance of 2,4-D Use on Soybeans: Total Ohio Acres Planted (4,850,000)
(see notes at end of table)

Year	Percent Acres Treated	Rate of Application (lbs/acre)	Number of Applications	Rate per Crop Year (lbs/acre)	Pounds Applied
2016	34.0	0.520	1.0	0.520	858,208
2015	34.0	0.520	1.0	0.520	840,513
2014	33.7	0.515	1.0	0.518	819,687
2013	33.3	0.511	1.0	0.518	776,993
2012	33.0	0.510	1.0	0.522	792,258
2011	32.3	0.508	1.0	0.518	762,117
2010	31.7	0.506	1.0	0.514	748,420
2009	31.0	0.505	1.0	0.511	720,857
2008	30.3	0.505	1.0	0.509	695,175
2007	29.7	0.506	1.0	0.507	639,582
2006	29.0	0.507	1.0	0.507	683,876
2005	19.0	0.421	0.8	0.422	360,383
2004	9.0	0.486	1.0	0.491	196,690
2003	6.5	0.422	0.8	0.426	119,003
2002	4.0	0.530	1.0	0.530	100,700
2001	4.0	0.320	1.0	0.320	58,880
2000	10.0	0.400	1.0	0.400	178,000
1999	9.0	0.570	1.0	0.580	240,120
1998	22.0	0.360	1.0	0.370	358,160
1997	9.0	0.440	1.0	0.440	172,260
1996	20.0	0.400	1.0	0.400	360,000
1995	15.0	0.470	1.0	0.470	285,525
1994	10.0	0.490	1.1	0.520	208,000
1993	6.0	0.520	1.0	0.520	129,480

- Notes:**
1. 2,4-D is sold in more than one chemical form, and surveyed separately by the USDA’s National Agricultural Statistics Service (NASS). Data on percent acres treated, number of acres treated, and pounds applied are the sum across all forms of the chemical. Rates of application and number of applications are averages across each form of the pesticide, weighted by shares of total acres treated.
 2. For years not surveyed by NASS, values are interpolated between the nearest two years with reported values. Values between the last survey and 2016 are extrapolated assuming no change in rate of application, number of applications, or percent acres treated.
 3. Each year when NASS surveys a crop, the agency strives to include 85% to 90% of acres planted. NASS surveyed acres at the national level are lower than total acres planted. For this table, NASS surveyed acres were 4,850,000.