

Table 3C.5 Reliance of Glyphosate Use on Soybeans: Total Wisconsin Acres Planted (1,960,000)
(see notes at end of table)

Year	Percent Acres Treated	Rate of Application (lbs/acre)	Number of Applications	Rate per Crop Year (lbs/acre)	Pounds Applied
2016	105.0	0.969	1.3	1.280	2,635,220
2015	105.0	0.969	1.3	1.280	2,527,660
2014	107.0	0.970	1.3	1.252	2,412,309
2013	109.0	0.962	1.3	1.215	2,093,181
2012	111.0	0.944	1.2	1.169	2,218,058
2011	106.7	0.869	1.2	1.086	1,877,024
2010	102.3	0.813	1.1	1.029	1,726,401
2009	98.0	0.777	1.1	1.000	1,597,987
2008	93.7	0.766	1.1	1.005	1,515,037
2007	89.3	0.783	1.2	1.046	1,307,801
2006	85.0	0.832	1.3	1.131	1,585,733
2005	81.0	0.626	1.1	0.865	1,128,079
2004	77.0	0.519	1.0	0.725	893,360
2003	73.0	0.525	1.1	0.731	917,379
2002	69.0	0.665	1.3	0.909	965,888
2001	63.5	0.697	1.2	0.878	892,360
2000	58.0	0.750	1.1	0.870	782,130
1999	41.0	0.717	1.3	0.937	518,630
1998	24.0	0.683	1.4	1.003	276,828
1997	7.0	0.650	1.6	1.070	77,896

- Notes:**
1. Glyphosate is sold in more than one chemical form, and surveyed separately by the USDA's National Agricultural Statistics Service (NASS). Data on percent acres treated, number of acres treated, and pounds applied are the sum across all forms of the chemical. Rates of application and number of applications are averages across each form of the pesticide, weighted by shares of total acres treated.
 2. For years not surveyed by NASS, values are interpolated between the nearest two years with reported values. Values between the last survey and 2016 are extrapolated assuming no change in rate of application, number of applications, or percent acres treated.
 3. Each year when NASS surveys a crop, the agency strives to include 85% to 90% of acres planted. NASS surveyed acres at the national level are lower than total acres planted. For this table, NASS surveyed acres were 1,960,000.