

Table 3C.8 Reliance of S-metolachlor Use on Corn: Total Indiana Acres Planted (5,600,000)
 (see notes at end of table)

Year	Percent Acres Treated	Rate of Application (lbs/acre)	Number of Applications	Rate per Crop Year (lbs/acre)	Pounds Applied
2016	30.0	0.997	1.1	1.122	1,884,960
2015	30.0	0.997	1.1	1.122	1,901,790
2014	30.0	0.997	1.1	1.122	1,985,940
2013	34.0	0.988	1.1	1.095	2,233,800
2012	38.0	0.980	1.1	1.067	2,534,125
2011	42.0	0.972	1.1	1.040	2,577,120
2010	46.0	0.963	1.1	1.012	2,746,568
2009	44.4	1.017	1.1	1.078	2,680,339
2008	42.8	1.071	1.1	1.144	2,790,902
2007	41.2	1.126	1.1	1.209	3,237,702
2006	39.6	1.180	1.1	1.275	2,776,950
2005	38.0	1.234	1.1	1.341	3,006,522
2004	30.0	1.277	1.1	1.331	2,276,010
2003	22.0	1.320	1.0	1.320	1,626,240
2002	22.0	1.230	1.0	1.230	1,461,240
2001	32.0	1.240	1.0	1.250	2,320,000
2000	27.0	1.250	1.0	1.250	1,923,750

- Notes:**
1. For years not surveyed by NASS, values are interpolated between the nearest two years with reported values. Values between the last survey and 2016 are extrapolated assuming no change in rate of application, number of applications, or percent acres treated.
 2. Each year when NASS surveys a crop, the agency strives to include 85% to 90% of acres planted. NASS surveyed acres at the national level are lower than total acres planted. For this table, NASS surveyed acres were 5,600,000.