

Table 3C.8 Reliance of S-metolachlor Use on Corn: Total Ohio Acres Planted (3,550,000)
 (see notes at end of table)

Year	Percent Acres Treated	Rate of Application (lbs/acre)	Number of Applications	Rate per Crop Year (lbs/acre)	Pounds Applied
2016	32.0	1.123	1.0	1.167	1,325,712
2015	32.0	1.123	1.0	1.167	1,325,712
2014	32.0	1.123	1.0	1.167	1,381,728
2013	30.3	1.175	1.0	1.228	1,448,733
2012	28.5	1.227	1.1	1.289	1,432,724
2011	26.8	1.279	1.1	1.350	1,227,825
2010	25.0	1.331	1.1	1.411	1,216,988
2009	26.4	1.311	1.1	1.375	1,216,050
2008	27.8	1.292	1.1	1.340	1,229,316
2007	29.2	1.272	1.0	1.304	1,465,957
2006	30.6	1.253	1.0	1.269	1,223,189
2005	32.0	1.233	1.0	1.233	1,361,232
2004	23.0	1.232	1.0	1.232	949,256
2003	14.0	1.230	1.0	1.230	568,260
2002	24.0	1.290	1.0	1.290	1,006,200
2001	24.0	1.230	1.0	1.230	1,003,680
2000	18.0	1.030	1.0	1.030	658,170

- Notes:**
1. For years not surveyed by NASS, values are interpolated between the nearest two years with reported values. Values between the last survey and 2016 are extrapolated assuming no change in rate of application, number of applications, or percent acres treated.
 2. Each year when NASS surveys a crop, the agency strives to include 85% to 90% of acres planted. NASS surveyed acres at the national level are lower than total acres planted. For this table, NASS surveyed acres were 3,550,000.