

**Table 3C.9 Reliance of S-metolachlor Use on Corn: Total Indiana Acres Planted**  
(see notes at end of table)

Year	Total Acres Planted	Percent Acres Treated	Rate of Application (lbs/acre)	Number of Applications	Rate per Crop Year (lbs/acre)	Pounds Applied
2016 *	5,600,000	30.0	0.940	1.10	1.016	1,706,880
2015	5,650,000	30.0	0.968	1.10	1.069	1,811,955
2014 *	5,900,000	30.0	0.997	1.10	1.122	1,985,940
2013	6,000,000	34.0	0.988	1.10	1.095	2,233,800
2012	6,250,000	38.0	0.980	1.10	1.067	2,534,125
2011	5,900,000	42.0	0.972	1.10	1.040	2,577,120
2010 *	5,900,000	46.0	0.963	1.10	1.012	2,746,568
2009	5,600,000	44.4	1.017	1.10	1.078	2,680,339
2008	5,700,000	42.8	1.071	1.10	1.144	2,790,902
2007	6,500,000	41.2	1.126	1.10	1.209	3,237,702
2006	5,500,000	39.6	1.180	1.10	1.275	2,776,950
2005 *	5,900,000	38.0	1.234	1.10	1.341	3,006,522
2004	5,700,000	30.0	1.277	1.05	1.331	2,276,010
2003 *	5,600,000	22.0	1.320	1.00	1.320	1,626,240
2002 *	5,400,000	22.0	1.230	1.00	1.230	1,461,240
2001 *	5,800,000	32.0	1.240	1.00	1.250	2,320,000
2000 *	5,700,000	27.0	1.250	1.00	1.250	1,923,750

**Notes:**

1. S-metolachlor is sold in more than one chemical form, and surveyed separately by the USDA's National Agricultural Statistics Service (NASS). Data on percent acres treated, number of acres treated, and pounds applied are the sum across all forms of the chemical. Rates of application and number of applications are averages across each form of the pesticide, weighted by shares of total acres treated.
2. For years not surveyed by NASS, values are interpolated between the nearest two years with reported values. An asterisk denotes which years were surveyed by NASS.
3. Each year when NASS surveys a crop, the agency strives to include 85% to 90% of acres planted. NASS surveyed acres at the national level are lower than total acres planted. The above data is indicative of pesticides applied to total acres planted.