

Table 3C.9 Reliance of S-metolachlor Use on Corn: Total Wisconsin Acres Planted
(see notes at end of table)

Year	Total Acres Planted	Percent Acres Treated	Rate of Application (lbs/acre)	Number of Applications	Rate per Crop Year (lbs/acre)	Pounds Applied
2016 *	4,050,000	28.0	1.275	1.10	1.350	1,530,900
2015	4,000,000	28.0	1.256	1.05	1.306	1,462,720
2014 *	4,000,000	28.0	1.237	1.00	1.263	1,414,560
2013	4,100,000	29.0	1.318	1.00	1.337	1,589,693
2012	4,350,000	30.0	1.398	1.00	1.411	1,841,355
2011	4,150,000	31.0	1.478	1.00	1.485	1,910,453
2010 *	3,900,000	32.0	1.559	1.00	1.559	1,945,632
2009	3,850,000	31.6	1.542	1.00	1.542	1,875,997
2008	3,800,000	31.2	1.525	1.00	1.525	1,808,040
2007	4,050,000	30.8	1.507	1.00	1.507	1,879,832
2006	3,650,000	30.4	1.490	1.00	1.490	1,653,304
2005 *	3,800,000	30.0	1.473	1.00	1.473	1,679,220
2004	3,600,000	29.0	1.512	1.00	1.512	1,578,528
2003 *	3,750,000	28.0	1.550	1.00	1.550	1,627,500
2002 *	3,650,000	10.0	1.600	1.00	1.600	584,000
2001 *	3,400,000	18.0	1.280	1.00	1.280	783,360
2000 *	3,500,000	15.0	1.180	1.00	1.190	624,750

Notes:

1. S-metolachlor is sold in more than one chemical form, and surveyed separately by the USDA's National Agricultural Statistics Service (NASS). Data on percent acres treated, number of acres treated, and pounds applied are the sum across all forms of the chemical. Rates of application and number of applications are averages across each form of the pesticide, weighted by shares of total acres treated.
2. For years not surveyed by NASS, values are interpolated between the nearest two years with reported values. An asterisk denotes which years were surveyed by NASS.
3. Each year when NASS surveys a crop, the agency strives to include 85% to 90% of acres planted. NASS surveyed acres at the national level are lower than total acres planted. The above data is indicative of pesticides applied to total acres planted.